



**Laboratory Guide to Massive Transfusion**

**Definition:** Anticipated loss of 90% or more of a patient's total blood volume in 3 hours or less and/or anticipated need for more than 10 units of blood in a 24 hour period.

**Early recognition**

**Prompt action**

**Good Communication**

Objective	Main Points	Comments
<b>Restore circulating volume</b>	<ol style="list-style-type: none"> <li>1. Insert wide-bore IV</li> <li>2. Give adequate volumes of warmed crystalloid (+/- colloid)</li> <li>3. Aim to maintain normal BP and urine output of &gt;30 ml/hr</li> </ol>	<ul style="list-style-type: none"> <li>-14 G or larger</li> <li>-Monitor CVP</li> <li>-Blood loss often underestimated</li> </ul>
<b>Initiate massive transfusion protocol</b>	<ol style="list-style-type: none"> <li>1. Designate one person to communicate with Transfusion Lab.</li> <li>2. Call Transfusion Lab to initiate massive transfusion protocol.</li> <li>3. Confirm the availability of crossmatch specimen and estimated delivery time for blood, plasma and platelets.</li> <li>4. Order crossmatch for 6 units of packed red cells.</li> <li>5. Consider unmatched or group specific red cells.</li> <li>6. Consider blood salvage if available and appropriate.</li> </ol>	<ul style="list-style-type: none"> <li>-Unmatched O Neg – 10 minutes</li> <li>-Unmatched ABO group specific – 15 minutes *</li> <li>-Fully crossmatched – 45 minutes*</li> <li>*Excluding collection and delivery time for crossmatch specimen. A stat crossmatch can be performed in 10 minutes if a pre-operative group and screen has been done and the patient does not have alloantibodies.</li> </ul>
<b>Transfusion Laboratory protocol activated</b>	<ol style="list-style-type: none"> <li>1. Notify Hematopathologist.</li> <li>2. Notify Hematology lab to optimize turn around time of stat blood work.</li> <li>3. Issue multiple units of red cells in the most expedient fashion possible.</li> <li>4. Assess platelet stores and order enough to have 10 units available.</li> <li>5. Thaw 2 units of FFP.</li> <li>6. Issue up to 6 units of FFP and one adult dose of platelets without requirement for supporting lab data or pathologist approval.</li> <li>7. Keep 6 units RBC crossmatched on hand.</li> </ol>	<ul style="list-style-type: none"> <li>-Transfusion lab staff empowered to issue blood components without waiting for lab data or pathologist approval.</li> </ul>
<b>Avoid hypothermia</b>	<ol style="list-style-type: none"> <li>1. Pre-warmed crystalloid</li> <li>2. Rapid infusion blood warmer</li> <li>3. Warm ambient room temp</li> <li>4. Warming/reflective blankets</li> <li>5. Warm saline for irrigation</li> <li>6. Warmed &amp; humidified anaesthetic gases</li> </ol>	<ul style="list-style-type: none"> <li>-Hypothermia impairs coagulation and platelet function.</li> <li>-Most under recognized cause of coagulopathy.</li> <li>-Common in massive transfusion.</li> <li>-Worse in thoracic / abdominal surgery.</li> <li>-Prophylactic FFP contributes to hypothermia.</li> <li>-Aim for temp &gt; 35°C.</li> </ul>
<b>Achieve hemostasis</b>	<ol style="list-style-type: none"> <li>1. Treat any surgical source of bleeding.</li> <li>2. Correct coagulopathy with judicious use of blood components.</li> </ol>	
<b>Order lab tests</b>	<ol style="list-style-type: none"> <li>1. CBC, INR, PTT, CBC, Fibrinogen after 6 units packed cells.</li> <li>2. Repeat as required to guide component therapy.</li> </ol>	
<b>Request platelets</b>	<ol style="list-style-type: none"> <li>1. Not always available on site.</li> <li>2. Allow 1-2 hours for delivery from blood centre.</li> <li>3. Expect count&lt;50 with 2 x blood volume replacement.</li> </ol>	<ul style="list-style-type: none"> <li>-Target: &gt;50 x 10<sup>9</sup>/L (&gt;100 is desirable for multiple or CNS trauma, however &gt;75 is more realistic)</li> <li>-Initial adult dose=1 buffy coat or 1 apheresis unit (Child &lt;20Kg: 10-15 mL/kg)</li> </ul>
<b>Request FFP</b>	<ol style="list-style-type: none"> <li>1. Consider after 6-10 units red cells or 1-2 x volume replacement.</li> <li>2. Ideally based upon INR/PTT results.</li> <li>3. Allow 30 minutes thawing time.</li> </ol>	<ul style="list-style-type: none"> <li>-Aim for PT and PTT &lt;1.5 x mid normal</li> <li>-Dosage 10 – 15 ml/kg (4 units/70 kg)</li> </ul>
<b>Request cryoprecipitate</b>	<ol style="list-style-type: none"> <li>1. Primarily to replace fibrinogen if &lt; 1.0 g/L.</li> <li>2. Expect &lt; 1 g/L with 1.5 x volume replacement.</li> <li>3. If time permits, measure fibrinogen first.</li> <li>4. Allow 30 minutes to thaw and pool.</li> </ol>	<ul style="list-style-type: none"> <li>-Dosage 1-1.5 units/10kg (10 units/70 kg)</li> <li>-Limited on site supply</li> <li>-For continued non-surgical bleeding despite plasma and platelet transfusion</li> </ul>

References: LA County & UCLA Trauma Surgery and Critical Care Protocols  
Stainsby, D.; MacLennan, S.; Hamilton, P.J. (2000). Management of massive blood loss: A template guideline. *British J of Anaesth*, 85(3): 487-91